

This is a repository copy of *Palliative care triggers in progressive neurodegenerative conditions : An evaluation using a multi-centre retrospective case record review and principal component analysis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/129071/>

Version: Accepted Version

Article:

Hussain, Jamilla, Allgar, Victoria Louise orcid.org/0000-0002-5228-2623 and Oliver, David (2018) Palliative care triggers in progressive neurodegenerative conditions : An evaluation using a multi-centre retrospective case record review and principal component analysis. Palliative Medicine. ISSN 0269-2163

<https://doi.org/10.1177/0269216318755884>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Palliative Medicine

Development of an evidence base for palliative care triggers in progressive neurodegenerative conditions: a multi-centre retrospective case record review and principal component analysis

Journal:	<i>Palliative Medicine</i>
Manuscript ID	PMJ-16-0358.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Hussain, Jamiila ; University of York, Hull York Medical School Allgar, Victoria; University of York, Hull York Medical School Oliver, David; University of Kent, Tizard Centre
Keywords:	Triggers, Progressive neurodegenerative conditions, Palliative care
Abstract:	<p>Background: The use of specific triggers has been suggested to help identify patients with progressive neurological disease who would benefit from palliative care.</p> <p>Aim: To develop an evidence base for the use of triggers for patients with progressive neurological disease.</p> <p>Design: A retrospective case note review was undertaken in 12 specialist palliative care units across the UK, extracting the timing, and presence of triggers in the last 2 years of life.</p> <p>Results: 300 records were reviewed: MND/ALS 58%, Parkinson's disease 17% and Parkinson's plus syndromes 12%. There was a high burden of triggers – 16 in the last 2 years of life and 10 in the last 6 months of life. Four factors were found to explain 64% of the total variance:</p> <p>Factor 1 – Deterioration in physical function, dysphagia, significant complex symptoms and pain</p> <p>Factor 2 – Weight loss and respiratory symptoms</p> <p>Factor 3 – Recurrent infections and cognitive decline</p> <p>Factor 4 – Aspiration pneumonia</p> <p>A Cox regression analysis showed a statistically significant association, reducing the hazard of death, between Factor 1 and survival. When all diagnoses were grouped together the total number of triggers at 3 months was associated with survival.</p> <p>Conclusion: This study shows that there are high burden of triggers in the last months and years of life and they may be valuable in predicting deterioration in the last 6 to 12 months of life for people with progressive neurological disease as the number of triggers and the frequency of triggers increases as death approaches.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

SCHOLARONE™
Manuscripts

For Peer Review

PALLIATIVE MEDICINE AUTHOR SUBMISSION CHECKLIST

Please complete this checklist for all papers submitted. Please answer questions with yes, no (and give reasons) or not applicable.

This checklist is a mandatory upload on submission, without it your paper will be returned without consideration.

Item	Explanation	Addressed? (Yes, No with reasons, Not applicable)
Article title	Have you followed our guidelines on writing a good title that will be found by search engines? (E.g. with methods in the title, use of common words for the issue addressed, no country names, and possibly indicating findings). If your study has an acronym is it included in the title?	Yes
Abstract	Have you followed our guidelines on writing your structured abstract? Please remember we have separate abstract structures for original research, reviews and case reports. There should be no abbreviations in the abstract, EXCEPT a study acronym which should be included if you have one. If a trial (or other design formally registered with a database) have you included your registration details?	Yes
Key statements	Have you included our key statements within the body of your paper (after abstract and before the main text is a good place!) and followed our guidelines for how these are to be written? There are three main headings required, and each may have 1-3 separate bullet points. Please use clear, succinct, single sentence separate bullet points rather than complex or multiple sentences.	Yes
Keywords	Have you given keywords for your study? We ask that these are current MeSH headings unless there is no suitable heading for use (please give explanation in cover letter).	Yes
International relevance	Have you contextualised your work for an international audience and explained how your work contributes to an international knowledge base?	Yes
Publishing guidelines	Have you submitted a checklist for a relevant publishing guideline as a supplementary file? These include CONSORT, PRISMA, COREQ checklists, but others may be more relevant for your type of manuscript. If no published checklist exists please create one as a table from the list of requirements in your chosen guideline. If your study design does not have a relevant publishing guideline please review closest	Not applicable

	matches and use the most appropriate with an explanation.	
Word count	Does your paper adhere to our word count for your article type? Please insert number of words in the box to the right. Remember that tables, figures, qualitative data extracts and references are not included in the word count.	Yes
Figures and tables and/or quotations	Have you adhered to our guidelines on the number of tables and figures for your article type? Data (e.g. quotations) for qualitative studies are not included in the word count, and we prefer that they are integrated into the text (e.g. not in a separate table).	Yes
Study registration	Where appropriate have you included details (including reference number, date of registration and URL) of study registration on a database e.g. trials or review database. If your study has a published protocol, is this referenced within the paper?	Not applicable
Other study publications?	If there are other publications from this study are these referenced within the body of the paper? Please do not reference papers in preparation or submitted, but in-press publications are acceptable.	No
Scales, measures or questionnaires	If your study primarily reports the development or testing of scales/measures or questionnaires have you included a copy of the instrument as a supplementary file?	Not applicable
Abbreviations	Have you removed all abbreviations from the text except for extremely well known, standard abbreviations, which should be spelt out in full first? We particularly discourage abbreviations for core concepts such as palliative or end of life care.	Yes
Research ethics and governance approvals for research involving human subjects	Have you given full details of ethics/governance/data protection approvals with reference numbers, full name of the committee(s) giving approval and the date of approval? If such approvals are not required have you made it explicit within the paper why they were not required. Are details of consent procedures clear in the paper?	Yes
Date(s) of data collection	Have you given the dates of data collection for your study within the body of your text? If your data are over 5 years old you will need to articulate clearly why they are still relevant and important to current practice.	Yes

Structured discussion	Does your paper contain a structured discussion summarising the main findings, addressing strengths and limitations, articulating what this study adds, and presenting the implications for practice.	Yes
Case reports	If your study is a case report have you followed our clear structure for a case report, including highlighting what research is needed to address the issue raised? Have you made clear what consent was required or given for the publication of the case report? Have you provided evidence of such consent as a supplementary file to the editor?	Not applicable
Acknowledgements and declarations	Have you included a funding declaration according to the SAGE format? Are there acknowledgements to be made? Have you stated where data from the study are deposited and how they may be available to others? Have you conflicts of interest to declare?	Not applicable
Supplementary data and materials	Is there any content which could be provided as supplementary data which would appear only in the online version of accepted papers? This could include large tables, full search strategies for reviews, additional data etc.	Yes
References	Are your references provided in SAGE Vancouver style?	Yes
Ownership of work.	Can you assert that you are submitting your original work, that you have the rights in the work, that you are submitting the work for first publication in the Journal and that it is not being considered for publication elsewhere and has not already been published elsewhere, and that you have obtained and can supply all necessary permissions for the reproduction of any copyright works not owned by you.	Yes

Palliative care triggers in progressive neurodegenerative conditions: a multi-centre retrospective case record review and principal component analysis

Jamilla Hussain
NIHR Doctoral Research Fellow,
Hull York Medical School
University of York,
York, YO10 5DD

Victoria Allgar
Senior Lecturer
HYMS/Health Sciences
University of York
York YO10 5DD

David Oliver
Honorary Professor
Tizard Centre
University of Kent
Canterbury

Corresponding author:

Professor David Oliver
Tizard Centre
Cornwallis North East
University of Kent
Canterbury
Kent CT2 7NF

Tel: 01634 828485
Email: drdjoliver@gmail.com

Abstract

Background: The use of specific triggers has been suggested to help identify patients with progressive neurological disease who would benefit from palliative care.

Aim: This study aimed to develop an evidence base for the use of triggers for patients with progressive neurological disease.

Design: A retrospective case note review of the timing and presence of triggers in the last 2 years of life was undertaken.

Setting/participants: 12 specialist palliative care units across the UK provided data from 300 patients: mean patient age 70 years, 50% male, diagnoses included Motor Neurone Disease 58%, Parkinson's disease 17% and Parkinson's plus syndromes 12%.

Results: There was a high burden of triggers – 17 in the last 2 years of life and 10 in the last 6 months of life. The most frequent triggers were: deteriorating physical function, complex symptoms and dysphagia. Four factors were found to explain 64% of the total variance:

Factor 1 – Deterioration in physical function, dysphagia, significant complex symptoms and pain

Factor 2 – Weight loss and respiratory symptoms

Factor 3 – Recurrent infections and cognitive decline

Factor 4 – Aspiration pneumonia.

Cox regression analyses found different triggers were associated with survival from diagnosis and referral to palliative care across all participants, and for different neurological conditions..

Conclusions: This study demonstrates that there is a high burden of triggers in the last months and years of life and that these could potentially be reduced to fewer components. Prospective studies assessing which triggers are useful for different conditions are now required.

Key Words

Triggers, palliative care, neurodegenerative disease, survival analysis, Motor Neuron Disease, Parkinson’s Disease

What is already known about the topic?

The use of triggers to identify the end of life phase and need for palliative care involvement for patients with progressive neurological conditions has been advocated in several policy documents.

The triggers suggested are based on expert consensus.

There is a need to build an evidence base to inform and evaluate such policy recommendations.

What this paper adds?

The average number of triggers in the last 2 years of life was 17, with an exponential increase in the last 6 months of life.

1 Four factors explained 64% of the variance in the triggers.

2

3 Different triggers were associated with survival from diagnosis and referral to
4 palliative care across all participants, and for different neurological conditions.

5

6 **Implications for practice, theory or policy?**

7 The high burden of triggers in the last few months of life indicates the need for
8 palliative care involvement for this patient group.

9
10 The correlation between triggers suggests the triggers could be reduced to fewer
11 components.

12
13 There is evidence that different triggers may help prognostication over different time
14 frames and for different conditions.

15

16

17

18

19

20

21

22

23

24

25

26

27

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

The care of people with neurological disease is complex and is a challenge for patients, families and professionals. There is a perceived need for palliative care for this patient group, in particular those with progressive degenerative neurological conditions (PNCs), such as motor neurone disease (MND), Multiple sclerosis (MS), Huntington’s disease (HD), Parkinson’s disease (PD) and the Parkinson Plus syndromes (PD Plus).

The need for palliative care has been discussed within specific disease guidelines (1,2) and for all neurological patients, and in the recent consensus document from the European Association for Palliative Care (EAPC) and the European Academy of Neurology (EAN) it was stressed that ‘palliative care should be considered early in the disease trajectory, depending on the underlying diagnosis’(3).

Although there is a perceived need, many patients with progressive neurological disease do not receive palliative care, and the access to specialist services is more

restricted than for cancer patients (4). It has been suggested that this is due to the variability in the progression and prognosis of patients with neurological disease and the difficulty in recognising deterioration and that a patient is at the end of life (4,5).

The National End of Life Care (NEoLC) Programme framework for end of life care in long term neurological conditions suggested possible triggers for the identification of the end of life phase (the last 6-12 months) in this patient population. These 'triggers' are characteristics or events which have a significance within the disease progression, are readily recognised and can be easily used clinically. The triggers suggested included swallowing problems, recurring infection, marked decline in functional status, first episode of aspiration pneumonia, cognitive difficulties, weight loss and significant complex symptoms (5). The use of such triggers has also been advocated by the Supportive and Palliative Care Indicators Tool (SPICT) (6) and the Marie Curie Triggers for Palliative Care (7) guidance which have suggested that triggers for palliative care involvement are used by service providers to improve palliative care access for such patients.

The triggers suggested by the NEoLC programme and other guidance are based on expert consensus and there has been little research in this area. It is essential that a robust evidence base is developed to inform and evaluate new palliative care policy, as was illustrated in the review of the Liverpool Care Pathway (LCP) (8). A small study from one centre evaluated the triggers suggested for PNCs and found four symptom components explained 76.8% of the variance (9). These triggers were rapid

1 physical decline, significant complex symptoms including pain, infection and
2 cognitive impairment, and risk of aspiration. In order to further assess the value of the
3 triggers, this study builds on this initial assessment and involved several centres in the
4 UK.

5
6 The objectives of this study were to explore (i) the frequency of triggers for palliative
7 care involvement in PNC in the last 2 years, and 6 months of life, and therefore to
8 identify which triggers are most burdensome for patients with PNCs; (ii) whether the
9 triggers were correlated and if the number of triggers could be reduced to fewer
10 components; (iii) the relationship between the triggers and trigger components, and
11 survival from diagnosis and referral to palliative care services and (iv) the
12 associations between triggers and survival for different diagnoses.

13
14 **Methods**

15
16 *Study design and setting*

17 A retrospective case-note review was conducted by 12 sites from across England and
18 Wales identified through the Association of Palliative Medicine Neurology Specialist
19 Interest Forum (APM Neuro-SIF). All were specialist palliative care services that
20 provide care at: home, hospice unit, in a day hospice or hospital. Data was extracted
21 between January 2014 and 2015. The study was discussed with Leeds East Health
22 Research Ethics Committee and it was agreed that the study met the criteria for UK
23 ethical regulations for research limited to secondary use of anonymised information

1 previously collected in the course of normal care and did not require review by the
2 research ethics committee.

4 *Participants*

5 Consecutive patients who had a diagnosis of a PNC, were under the care of a
6 specialist palliative care service, and had died between January 2009 and 2014 were
7 eligible for inclusion. The local site identified all participants.

9 *Data sources*

10 Members of the clinical team extracted data from the patients' clinical records,
11 including paper case-notes from hospices, hospitals and community teams, as well as
12 electronic databases. A standardised data collection form and a data collection
13 guideline were used to ensure consistency of data extraction across all centres. All
14 data were anonymised locally.

16 *Variables*

17 Data collected included demographic details (e.g. age, gender, and ethnicity), medical
18 history (diagnosis, date of diagnosis, number of comorbidities, date of palliative care
19 referral). For each trigger, according to documented evidence, the timing (in units of
20 months prior to death) of the first presentation and subsequent deterioration of the
21 trigger over the last two years of life were extracted. In addition to the triggers
22 assessed by Hussain et al (9), respiratory symptoms was also included as a trigger as

1 this was considered by the APM Neuro-SIF as a potential important trigger for
2 palliative care involvement in PNC, and in particular MND.
3
4
5
6
7
8
9 *Analysis*
10 Descriptive data is summarised by the mean (standard deviation (SD)) or number (%).
11
12 Principal component analysis (PCA) was used to assess the correlation between the
13 triggers and to identify common components and therefore whether the triggers could
14 be reduced to fewer components. In this type of analysis the correlation between
15 parameters (triggers) is attributed to their common dependence on independent
16 entities called ‘factors’. The coefficients that link parameters to factors are called
17 ‘factor loadings’; the number of factors is chosen to be as small as possible but large
18 enough to account for most of the variation within the data. PCA was conducted using
19 data from the last six months as there was evidence that the number of triggers
20 increased rapidly after this point and to optimise the number of complete cases. It was
21 decided *a-priori* that the number of factors in the varimax rotation would be based on
22 the number of eigenvalues >1.0 in the PCA. We adopted one common and
23 conventional rule of thumb to consider ‘factor loadings’ of 0.40 or larger to be ‘high’.

18 Tests of multicollinearity (Bartlett’s test of sphericity $p<0.05$) and sampling adequacy
(Kaiser-Meyer-Olkin (KMO)) were undertaken to check the analysis was appropriate.

20 The internal consistency and reliability of the factors was assessed using Cronbach’s
21 α . The least squares regression approach was used to calculate the factor scores,
22 which are standardized to a mean of zero. The factor scores may then be used as
23 variables in subsequent modelling (10).

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
- 1
 - 2 Univariable and multivariable (adjusted for age, gender, diagnosis and comorbidities)
 - 3 Cox regression analyses were used to assess the association of survival from (i)
 - 4 diagnosis and (ii) referral, and:
 - 5 a. factor scores determined by the PCA,
 - 6 b. individual triggers and the number of triggers at 3, 6 and 12 months.

The Cox regression analyses for survival from diagnosis were repeated according to whether the participants were diagnosed with MND, PD or PD plus. Assessment of the other diagnoses was not possible as the number of participants with the diagnoses was insufficient. A p-value of <0.05 was considered to indicate statistical significance and all analyses were undertaken on STATA (v14.0)

Results

In total 300 clinical records were reviewed retrospectively. The mean age was 70 years (range 35 to 98), 50% were male and 92% were White. The main diagnoses were MND (58%), PD (17%), PD Plus (12%), MS (9%), and HD (2%). The majority had co-morbidities (76%), with 46% having two or more comorbidities.

Frequency of triggers for palliative care involvement in PNC in the last 2 years and 6 months of life

1 The average number of total triggers over the last 2 years of life was 16.7 (SD 12.7)
2 and in the last 6 months 10.0 (SD 7.4). Table 1 demonstrates that the most frequent
3 triggers in both the last 2 years and 6 months of life was deteriorating physical
4 function (2 years: 5.0 (SD 4.0), 6 months: 2.9 (SD 2.5)) followed by significant
5 complex symptoms (2 years: 3.9 (SD 5.2), 6 months: 2.3 (SD 3.1)) and dysphagia (2
6 years: 2.5 (SD 2.3), 6 months: 1.5 (SD 1.5)). The same pattern was seen when trigger
7 frequency was assessed per diagnosis for patients with MND, PD, and PD Plus
8 (Appendix Table 1). However for those with MS significant complex symptoms (3.0
9 (SD 3.5)) were slightly more frequent than deteriorating physical function (2.8 (SD
10 3.1)), and for the seven participants diagnosed with Huntington’s disease cognitive
11 impairment was the second most frequent trigger (2.0 (SD 2.1)).

12
13 In the last 2 years of life just over half of patients had documented evidence of weight
14 loss (56%) and pain (56%), and in the last 6 months the figures were just under half
15 (weight loss 42%; pain 44%). Figure 1 illustrates the trend in the total number of
16 triggers according to months prior to death, this demonstrates an exponential increase
17 in the number of triggers over time, with a rapid increase in the last 6 months of life. .

18
19 **Principal component analysis**

20 Factor analysis yielded four separate factors that explained 64% of the total variance
21 in the data set when the eigenvalue =1 criterion was used. Using data from the last 6
22 months of life, the correlation matrix is shown in Table 2. Bartlett’s test of sphericity
23 ($\chi^2=349$, degree of freedom=36, $p<0.0001$) indicated that the correlation between the

variables were sufficiently large for PCA. The KMO measure of sampling adequacy was 0.68, which can be interpreted as the degree of common variance among the variables, and verified the sampling adequacy of the analysis.

Factor analysis, derived from the factor loadings and the analysis of the triggers in last 6 months of life, identified 4 factors, with the following groupings (Table 3):

- Factor 1. Deterioration in physical function, dysphagia, significant complex symptoms and pain
- Factor 2. Weight loss and respiratory symptoms
- Factor 3. Recurrent infections and cognitive decline
- Factor 4. Aspiration.

Factor 1 explained 22% of the variance, the second factor 16%, third factor 14%, and the fourth 12%. Factors 2-4 only loaded on one or two items, so must be interpreted with caution. Cronbach's α were 0.67 for factor 1, 0.37 for factor 2 and 0.26 for factor 3. If "respiratory symptoms" is excluded as a trigger then there is evidence that weight loss, cognitive impairment and infections cluster on the same component and explained 16% of the total variance (data not shown). Higher factor scores on each factor were associated with a higher number of triggers.

Association between triggers and survival in patients with PNC

1 The mean survival time from diagnosis was 56.5 (SD 6.3) months (95% CI: 43.3,
2 67.8) and the median survival 24 months (interquartile range: 64, 11).. The mean
3 survival from referral to palliative care was 29.1 (SD 6.0) months (95% CI: 17.4,
4 40.9) and the median survival was 10 months (interquartile range: 24, 3))..

5
6 *Survival and factor scores:*

7 In a Cox regression analysis assessing the association between the factor scores and
8 survival from diagnosis, there was no statistically significant association at the 5%
9 significant level. This remained the case following adjustments for age, gender,
10 diagnosis and number of co-morbidities (Table 4). In the Cox regression analysis
11 assessing the association between the factor scores and survival from palliative care
12 referral, factor 1 had a statistically significant association (hazard ratio (HR) 0.9, 95%
13 CI 0.76, 0.99)), this remained the case after adjusting for age, gender, diagnosis and
14 number of co-morbidities (HR 0.86, 95% CI 0.75, 0.99). The hazard ratio indicates
15 that an increase in factor 1 scores (deterioration in physical function, dysphagia,
16 significant complex symptoms and pain) reduces the risk of death, after adjustment
17 for the effects of the other variables in the model.

18
19 *Survival and individual triggers:*

20 In a multivariable Cox regression analysis assessing the association between
21 individual triggers and survival from diagnosis, there was evidence that the number of
22 triggers 3 months prior to death and diagnosis had a statistically significant
23 association with survival from diagnosis (Appendix Table 2). When repeated for

1 survival from referral to palliative care, there was evidence that again the number of
2 triggers 3 months prior to death had a statistically significant association with
3 survival, as well as weight-loss in the last 2 years of life and age. There was evidence
4 that the number of documented episodes of cognitive impairment was also associated
5 with survival, however the 95% CI for this crosses 1. As age, episodes of weight-loss,
6 and number of triggers at 3 months increased, the hazard of death increased
7 (Appendix Table 2).

8

9 **Association between triggers and survival for different diagnoses**

10 *Survival and factor scores:*

11 The multivariable Cox regression analyses for survival from diagnosis for the MND,
12 PD and PD Plus groups, found evidence that different factors were associated with
13 survival for the different diagnoses (Table 5). For MND there was insufficient
14 evidence that any of the factors scores were associated with survival when age, gender
15 and comorbidities were taken into account. For PD, there was evidence of a
16 significant association with factor 4 (aspiration), and for PD plus there was evidence
17 of a significant association with factor 1 (deterioration in physical function, dysphagia,
18 significant complex symptoms and pain) and factor 3 (recurrent infections and
19 cognitive decline).

20
21
22 The multivariable Cox regression analyses for survival from palliative care referral
23 for the MND, PD and PD Plus groups, also found evidence that different factors were

1 associated with survival for the different diagnoses (Table 5). For MND there was
2 insufficient evidence that any of the factors scores were associated with survival when
3 age, gender and comorbidities were taken into account. For PD, there was evidence of
4 a significant association with factor 1 and factor 4, and for PD plus there was
5 evidence of a significant association with factor 2, factor 3 and factor 4.

6
7 *Survival and individual triggers:*

8 The multivariable Cox regression analysis assessing the association between
9 individual triggers and survival from diagnosis, found for MND: being female and the
10 number of triggers at 3 months prior to death increased the hazard of death, for PD:
11 increasing number of aspirations increased the hazard of death, for PD plus: more
12 infections and episodes of cognitive impairment increased the hazard of death,
13 however as the total number of triggers at 6 months increased the hazard of death was
14 found to decrease (Appendix Table 3).

15
16 **Discussion**

17
18 This is the largest study to date to assess the value of the triggers for palliative care
19 involvement proposed by current palliative care policy guidance for individuals with
20 PNCs. There was evidence of high burden of triggers in the last 2 years of life for
21 patients with neurological conditions, with a rapid increase in the last 6 months. Four
22 factors explained a large proportion of the variance in the triggers indicating the
23 triggers could be grouped in fewer components. Different factors and triggers were

1 associated with survival from diagnosis and referral to palliative care services; this
2 was also the case when the associations were assessed for individual diagnoses. These
3 results indicate that the association of the triggers with survival is complex and that
4 different triggers may be more important in different PNCs.

5
6 The use of triggers appears to be valuable in predicting deterioration in the last 6 to 12
7 months of life. The commonest triggers were decline in physical function, complex
8 symptoms and dysphagia. These were found for all diagnoses. The frequency of these
9 triggers may reflect that they are the most burdensome issues for patients, but may
10 also reflect that the other triggers, especially weight loss and cognitive impairment are
11 less well assessed or documented. The total number of triggers increased as death
12 approached and there appears to be an exponential pattern as shown in the earlier
13 study (9). There is a rapid change in the numbers of triggers towards death and thus
14 monitoring the rate of change in the total number of triggers may be a useful
15 prognostic tool, indicating that the person may be in the last few months of life.

16
17 The four factors that explained the most variance in the last 6 months of life could be
18 categorised as factor 1: deterioration of physical function (which would include
19 deterioration in swallowing ability, development of significant complex symptoms,
20 and pain), factor 2: weight loss and respiratory symptoms, factor 3: recurrent
21 infections and cognitive impairment, and factor 4: aspiration. These are similar to the
22 factors identified in the previous analysis on a smaller sample (9). Despite the larger
23 sample in this study, the small Cronbach alphas indicate that the results should be

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 interpreted with caution. Current clinical understanding may not necessarily suggest
2 these groupings, and initially the expert group did report these triggers individually.
3 However the analyses in both studies have suggested similar factors and the
4 components do measure a large proportion of the variation in the data and therefore it
5 would seem that the triggers could be reduced to fewer components. Further studies
6 that collect the triggers data prospectively are now needed to reassess the factor
7 loading before implementing changes to practice.

8
9 The association between triggers and survival was complex. There was insufficient
10 evidence that the factors derived from the PCA were associated with survival from
11 diagnosis when all diagnoses were grouped together, however in the PD group factor
12 4 (aspiration) had a statistically significant association with survival from diagnosis
13 and for the PD plus group both factors 1 and 3. In terms of the individual triggers,
14 there was evidence when all diagnoses are grouped together that the total number of
15 triggers at 3 months was associated with survival from both diagnosis and palliative
16 care referral. Although there was evidence that as the number of episodes of weight
17 loss increased, the hazard of death increased when all the diagnoses were grouped
18 together, for the MND, PD and PD plus group other triggers had a significant
19 association with survival. This indicates that different triggers may be useful in aiding
20 prognostication for different conditions, however further research with larger samples
21 for each diagnostic group is required.

22
23 Limitations

1 Data collected as part of routine clinical practice was extracted for this study therefore
2 there is a risk of information bias due to inaccurate collection, interpretation or
3 documentation of triggers. To minimise bias at the data extraction phase detailed
4 guidance was provided, together with email / telephone support throughout the
5 process. In addition healthcare professionals who were aware of the clinical context
6 extracted data. This review focussed on triggers in the last two years of life and was
7 limited to patients known to palliative care services, thus these findings may not be
8 generalisable to all patients with PNCs. This should be addressed in future research..
9
10 This study provides evidence that the triggers for palliative care involvement
11 advocated by the NEoLCP, SPICT and Marie Curie Triggers for palliative care
12 guidance may be helpful in the assessment of patients with PNC and identifying
13 patients in the last few months of life. There is increasing evidence that palliative care
14 can be helpful in improving symptoms and quality of life (15, 16,17) and that the
15 involvement of SPC should be dependent on need rather than prognosis. However this
16 episodic approach is a challenge for SPC services (4, 18) and the use of triggers may
17 be able to help in both the identification of disease burden and prognostication that
18 death may be approaching, which would support the involvement of palliative care
19 with patients with PNC. It is essential however that the triggers suggested by experts
20 in the field are rigorously assessed and developed to ensure patients receive optimal
21 palliative care input. This study has helped to identify key areas for further
22 prospective research including how the triggers could potentially be categorised into
23 fewer components, how the rate of change of triggers is associated with survival and

1 which triggers are most useful for different PNC. We would also advocate patient and
2 carer involvement to determine which triggers for palliative care involvement warrant
3 further assessment.
4
5

For Peer Review

Acknowledgements

The authors would like to acknowledge the Centres who took part in this study:

Dove House Hospice, Hull - Dr Rachael Dixon

Douglas Macmillan Hospice, Stoke-on-Trent - Dr Caroline Bruckner-Holt

George Thomas Hospice and Marie Curie Hospice, Wales - Dr Siwan Seaman

Leicestershire and Rutland Hospice, Leicester - Professor Christina Faull

St. Catherine's Hospice, Scarborough - Debi Adams

St Clare Hospice, Hastingwood - Emily Stowe

St.Gemma's Hospice, Leeds - Dr Sophie Thomas

St. Leonard's Hospice, York - Dr Sarah Wilcox, Grace Duffy

Sue Ryder Nettlebed Hospice, Oxfordshire - Dr Pat Strubbe, Dr Katrien Naessens

Wheatfield's Hospice, Leeds - Dr Annette Edwards

Wisdom Hospice, Rochester - Professor David Oliver

Declaration of conflicting interests

The authors have no conflict of interest to declare

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

References

1. Miller RG, Rosenberg JA, Gelinas DF, et al. Practice parameter: the care of the patient with amyotrophic lateral sclerosis (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology: ALS Practice Parameters Task Force. *Neurology* 1999;**52**(7):1311-23.

2. Andersen PM, Abrahams S, Borasio GD, de Carvalho M, Chio A, Van Damme P, Hardiman O, Kollewe K, Morrison KE, Petri S, Pradat PF, Silani V, Tomik B, Wasner M, Weber M. The EFNS Task Force on Diagnosis and Management of Amyotrophic Lateral Sclerosis. EFNS guidelines on the Clinical Management of Amyotrophic Lateral Sclerosis (MALS) - revised report of an EFNS task force. *Eur J Neurol* 2012; 19: 360-75.

3. Oliver D, Borasio GD, Caraceni A, de Visser M, Grisold W, Lorenzl S, Veronese S, Voltz R A consensus review on the development of palliative care for patients with chronic and progressive neurological disease – *Eur J Neurol*. 2015 DOI:10.1111/ene.1288

4. Oliver D J. Palliative care for people with progressive neurological disease: what is the role? *JPall Care* 2014; 30: 298-301

5. National End of Life Care Programme. End of life care in long term neurological conditions: a framework for implementation. 2010.

6. Supportive and Palliative Care Indicators Tool (SPICT)
<http://www.spict.org.uk/the-spict>

7. Marie Curie . Triggers for palliative care. Improving access to care for people with diseases other than cancer. 2015.
<https://www.mariecurie.org.uk/globalassets/media/documents/policy/policy-publications/june-2015/triggers-for-palliative-care-full-report.pdf>

8. UK Department of Health. More Care, Less pathway. A Review of the Liverpool Care Pathway. 2013.
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/212450/Liverpool_Care_Pathway.pdf

9. Hussain J, Adams D, Allgar V, Campbell C. Triggers in advanced neurological conditions: prediction and management of the terminal phase. *BMJ Supp Pall Care* 2014; 4: 30-37.

10. DiStefano, Christine, Zhu, Min & Mîndrilă, Diana (2009). Understanding and Using Factor Scores: Considerations for the Applied Researcher. *Practical Assessment, Research & Evaluation*, 14 (20).

11. Association for Palliative Medicine of Great Britain and Ireland. Commissioning Guidance for Specialist Palliative Care: Helping to deliver commissioning objectives. 2012
<http://www.ncpc.org.uk/sites/default/files/CommissioningGuidanceforSpecialistPalliativeCare.pdf>

12. Public Health England. End of Life Care Profiles
<http://fingertips.phe.org.uk/profile/end-of-life/data#page/0/gid/1938132883/pat/6/par/E12000008/ati/102/are/E0600003> ,

13. Brogaard T, Neergaard MA, Sokolowski I, Olesen F, Jensen AB. Congruence between preferred and actual place of care and death among Danish cancer patients. *Pall Med* 2012; 27: 155-162.
14. National End of Life Care Intelligence Network. What do we know now that we didn't know a year ago. 2012
15. Veronese S, Gallo G, Valle A, Cugno C, Chio A, Calvo A, Cavalla P, Zibetti M, Rivoiro C, Oliver DJ. Specialist palliative care improves the quality of life in advanced neurodegenerative disorders: Ne-PAL, a pilot randomized controlled study. *BMJ Supp and Pall Care* 2015;):1-9. Doi:10.1136/bmjspcare-2014-000788
16. Edmonds P, Hart S, Gao W, Vivat B, Burman R, Silber E, Higginson I. Palliative care for people severely affected by multiple sclerosis: evaluation of a novel palliative care service. *Mult Scler* 2010; 16: 627-36.
17. Higginson IJ, et al., Is Short-Term Palliative Care Cost-Effective in Multiple Sclerosis? A Randomized Phase II Trial. *Journal of Pain and Symptom Management*, 2009; 38: 816-826.
18. Oliver D, Silber E. End of life care in neurological disease. In oliver D (ed) *End of Life Care in Neurological Disease*. 2013. London. Springer. Page 24

Table 1. Average number of each trigger in the last 2 years and 6 months of life and the proportion of patients with each trigger

Triggers	Average number of triggers in last 2 years of life Mean (SD)	Proportion of participants with the trigger in last 2 years of life Number (%)	Average number of triggers in last 6 months of life Mean (SD)	Proportion of participants with the trigger in last 6 months of life Number (%)
Deteriorating physical function	5.0 (4.0)	286 (96%)	2.9 (2.5)	267 (89%)
Dysphagia	2.5 (2.3)	258 (86%)	1.5 (1.5)	211 (70%)
Aspiration	0.8 (1.5)	119 (40%)	0.7 (0.1)	109 (36%)
Infection	0.8 (1.3)	123 (41%)	0.5 (0.9)	93 (31%)
Weight loss	1.2 (1.6)	168 (56%)	0.6 (0.9)	127 (42%)
Cognitive impairment	0.9 (2.1)	96 (32%)	0.5 (1.1)	75 (25%)
Significant complex symptoms	3.9 (5.2)	206 (69%)	2.3 (3.1)	182 (61%)
Pain	1.3 (2.3)	167 (56%)	0.8 (1.4)	133 (44%)
Respiratory symptoms	(1.0)	63 (21%)	0.3 (0.8)	42 (14%)

Table 2. Correlation matrix for each trigger in the last 6 months of life

	Decline in physical function	Dysphagia	Aspiration	Infection	Weight loss	Cognitive impairment
Decline in physical function	1					
Dysphagia	0.47 <0.001	1				
Aspiration	0.04 0.5	0.21 <0.001	1			
Infection	0.15 <0.01	0.01 0.9	-0.00 0.9	1		
Weight loss	0.32 <0.001	0.31 <0.001	0.02 0.8	0.09 0.1	1	
Cognitive impairment	0.08 0.2	0.11 0.06	-0.05 0.4	0.15 <0.01	0.07 0.2	1
Pain	0.29 <0.001	0.24 <0.001	0.13 <0.01	-0.05 0.4	0.01 0.8	0.01 0.9
Significant complex symptoms	0.54 <0.001	0.30 <0.001	0.01 0.8	0.09 0.1	0.19 0.001	0.06 0.3
Respiratory	0.19 0.001	0.20 <0.001	0.06 0.3	-0.04 0.5	0.23 <0.001	-0.07 0.2

Table 3. Summary of the principal component analysis using triggers in the last 6 months of life.

	Factor 1: Deterioration in physical function, dysphagia, significant complex symptoms and pain	Factor 2: Weight loss and respiratory symptoms	Factor 3: Recurrent infections and cognitive decline	
Pain	.76			
Significant complex symptoms	.74			
Decline in physical function	.73			
Dysphagia	.49			
Weight loss		.72		
Respiratory		.72		
Cognitive impairment			.73	
Infection			.71	
Aspiration				

Table 5. Cox regression analysis assessing the association of the factors derived from the principal component analysis with survival from diagnosis

Survival time	Explanatory variable	Motor neurone disease			Parkinson's disease			Parkin	
		Adjusted hazard ratio*	95% CI	P-value	Adjusted hazard ratio*	95% CI	P-value	Adjusted hazard ratio*	95%
From diagnosis	Factor 1	0.96	0.84, 1.09	0.5	0.79	0.49, 1.26	0.3	2.16	1.1
	Factor 2	1.04	0.90, 1.21	0.6	0.83	0.41, 1.70	0.6	0.86	0.5
	Factor 3	1.01	0.86, 1.18	0.9	1.30	0.83, 2.03	0.3	1.70	1.0
	Factor 4	0.97	0.81, 1.13	0.7	2.51	1.19, 5.32	<0.05	1.26	0.8
From palliative care referral	Factor 1	0.94	0.80, 1.12	0.5	0.58	0.33, 0.99	<0.05	1.38	0.6
	Factor 2	1.05	0.90, 1.22	0.6	0.63	0.36, 1.57	0.3	0.50	0.2
	Factor 3	1.00	0.85, 1.19	0.9	1.02	0.64, 1.64	0.9	2.49	1.3
	Factor 4	0.88	0.73, 1.05	0.2	2.47	1.06, 5.75	<0.05	2.06	1.2


* Model adjusted for age, gender and number of comorbidities

Table 4.

Survival time	Explanatory variable	Hazard ratio	95% CI	P-value	Adjusted hazard ratio*	95% CI	P-value
From diagnosis	Factor 1	1.03	0.93, 1.15	0.5	0.96	0.86, 1.07	0.5
	Factor 2	1.11	0.99, 1.26	0.1	1.00	0.88, 1.15	0.9
	Factor 3	0.96	0.85, 1.09	0.6	1.05	0.92, 1.19	0.5
	Factor 4	1.06	0.94, 1.20	0.3	1.01	0.88, 1.14	0.9
From palliative care referral	Factor 1	0.87	0.76, 0.99	<0.05	0.86	0.75, 0.99	<0.05
	Factor 2	1.05	0.93, 1.19	0.4	1.03	0.89, 1.18	0.7
	Factor 3	0.99	0.86, 1.13	0.9	1.04	0.90, 1.20	0.6
	Factor 4	1.03	0.90, 1.19	0.7	0.99	0.85, 1.16	0.9

* Model adjusted for age, gender, diagnosis and number of comorbidities

Figure 1: Total number of triggers (any) by month (Mean and standard error)



Appendix Table 1: Mean number of triggers in the last 2 years of life per diagnosis (SD)

Triggers	MND ¹ N=173	PD ² N=50	PD Plus ³ N=36	MS ⁴ N=26	HD ⁵ N=7
Deteriorating physical function	5.2 (4.0)	5.2 (4.0)	6.4 (4.5)	2.8 (3.1)	2.3 (1.8)
Dysphagia	2.7 (2.3)	1.8 (1.4)	3.4 (3.3)	1.5 (1.8)	1.4 (0.8)
Aspiration	0.8 (1.3)	0.7 (1.3)	1.3 (2.3)	0.8 (1.0)	1.3 (2.0)
Infection	0.7 (1.2)	0.7 (1.3)	1.4 (1.6)	1.3 (1.4)	0.6 (1.1)
Weight loss	1.2 (1.4)	1.5 (2.2)	1.3 (1.6)	0.7 (1.7)	1 (0.8)
Cognitive impairment	0.6 (2.1)	1.5 (2.0)	1.0 (1.7)	1.2 (2.0)	2.0 (2.1)
Significant complex symptoms	4.1 (5.0)	2.8 (3.1)	5.6 (8.5)	3.0 (3.5)	1.6 (2.1)
Pain	1.5 (2.6)	1.0 (1.4)	1.5 (2.0)	1.4 (2.4)	0.3 (0.5)
Respiratory symptoms	0.5 (1.2)	0.2 (0.7)	0.06 (0.2)	0.08 (0.3)	0

¹ Motor Neurone Disease

² Parkinson’s disease

³ Parkinson Plus syndromes

⁴ Multiple sclerosis

⁵ Huntington’s disease

Appendix Table 2: Cox regression analysis assessing the association of individual triggers and the total number of triggers at 3, 6 and 12 months with (i) survival from diagnosis and (ii) survival from palliative care referral (all participants)

Survival time	Explanatory variable	Univariable hazard ratio	95% CI	P-value	Adjusted hazard ratio*	95% CI	P-value
From diagnosis	Age	0.997	0.987, 1.006	0.5	1.006	0.993, 1.019	0.4
	Gender	0.836	0.655, 1.067	0.2	1.100	0.833, 1.454	0.5
	Diagnosis			<0.0001			<0.0001
	Comorbidities	1.025	0.942, 1.115	0.6	1.081	0.988, 1.183	0.09
	Decline in physical function	1.013	0.987, 1.039	0.4	0.977	0.927, 1.029	0.4
	Dysphagia	1.069	1.014, 1.127	0.01	1.046	0.964, 1.136	0.3
	Aspiration	1.031	0.953, 1.114	0.4	0.968	0.883, 1.062	0.5
	Infection	0.971	0.884, 1.066	0.5	0.977	0.873, 1.093	0.7
	Weight loss	1.030	0.960, 1.104	0.4	1.012	0.911, 1.124	0.8
	Cognitive impairment	0.974	0.913, 1.038	0.4	1.001	0.939, 1.068	1.0
	Pain	1.016	0.964, 1.070	0.6	0.990	0.928, 1.057	0.8
	Significant complex symptoms	1.012	0.990, 1.035	0.3	0.998	0.960, 1.036	0.9
	Respiratory symptoms	1.093	0.964, 1.239	0.2	0.894	0.749, 1.067	0.2
	Total number of triggers	1.006	0.997, 1.015		-	-	-
	Number of triggers at 3 months	1.078	1.021, 1.139	<0.01	1.083	1.013, 1.158	<0.05
	Number of triggers at 6 months	1.042	0.971, 1.118	0.3	1.028	0.934, 1.132	0.6
	Number of triggers at 12 months	0.969	0.866, 1.085	0.6	0.973	0.851, 1.113	0.7
From palliative care	Age	1.014	1.003, 1.025	<0.05	1.019	1.005, 1.034	<0.05
	Gender	1.186	0.924, 1.521	0.2	1.109	0.834, 1.484	0.5

referral			1.524			1.475	
	Diagnosis			0.7			0.5
	Decline in physical function	0.988	0.958, 1.019	0.4	0.954	0.904, 1.007	0.09
	Dysphagia	1.024	0.972, 1.078	0.4	1.005	0.932, 1.084	0.9
	Aspiration	1.005	0.913, 1.107	0.9	0.993	0.895, 1.101	0.9
	Infection	1.046	0.943, 1.160	0.4	1.115	0.988, 1.259	0.08
	Weight loss	1.049	0.974, 1.130	0.2	1.125	1.015, 1.246	<0.05
	Cognitive impairment	0.954	0.899, 1.013	0.1	0.933	0.870, 1.001	<0.05
	Pain	0.961	0.907, 1.017	0.2	0.981	0.921, 1.045	0.6
	Significant complex symptoms	0.993	0.967, 1.021	0.6	1.012	0.973, 1.052	0.6
	Respiratory symptoms	1.100	0.963, 1.257	0.2	0.999	0.841, 1.187	1.0
	Total number of triggers	0.997	0.987, 1.007	0.5	-	-	-
	Number of triggers at 3 months	1.052	0.994, 1.114	0.08	1.099	1.030, 1.171	<0.01
	Number of triggers at 6 months	1.002	0.923, 1.087	1.0	0.977	0.879, 1.086	0.7
	Number of triggers at 12 months	0.895	0.800, 1.002	0.1	0.912	0.799, 1.041	0.2

* Model adjusted for age, gender, diagnosis and number of comorbidities

Appendix Table 3: Multivariable Cox regression for survival from diagnosis and triggers for patients with Motor Neuron Disease, Parkinson's Disease and Parkinson Plus syndromes

Explanatory variable	Motor neurone disease		Parkinson's disease		Parkinson's plus	
	Hazard ratio	95% CI	Hazard ratio	95% CI	Hazard ratio	95% CI
Age	1.006	0.991, 1.021	1.058	0.989, 1.131	0.984	0.860, 1.125
Gender	1.422	1.003, 2.015	0.366	0.131, 1.019	0.591	0.159, 2.197
Comorbidities	1.082	0.963, 1.216	1.143	0.880, 1.484	1.053	0.716, 1.550
Decline in physical function	0.973	0.911, 1.040	1.081	0.885, 1.321	0.964	0.725, 1.281
Dysphagia	1.045	0.944, 1.156	1.017	0.721, 1.435	1.081	0.756, 1.547
Aspiration	0.907	0.794, 1.036	2.079	1.314, 3.289	1.149	0.914, 1.444
Infection	0.862	0.733, 1.012	1.195	0.892, 1.601	1.727	1.035, 2.916
Weight loss	1.084	0.936, 1.257	0.826	0.633, 1.658	1.157	0.653, 2.051
Cognitive impairment	0.993	0.919, 1.073	1.210	0.883, 1.658	1.515	1.024, 2.241
Pain	0.957	0.881, 1.040	1.123	0.772, 1.635	1.078	0.645, 1.801
Significant complex symptoms	0.977	0.928, 1.028	0.831	0.673, 1.027	1.123	0.986, 1.279
Respiratory symptoms	0.905	0.743, 1.102	1.416	0.616, 3.252	0.840	0.003, 218.2
Number of triggers at 3 months	1.129	1.044, 1.222	1.289	0.985, 1.688	1.131	0.794, 1.613
Number of triggers at 6 months	1.059	0.938, 1.196	0.826	0.606, 1.127	0.511	0.272, 0.963
Number of triggers at 12 months	0.930	0.759, 1.140	0.663	0.383, 1.147	1.338	0.787, 2.274